

The Improvement of the Bound on Hash Family

Xianmin Feng, Jiansheng Yang*

School of Science, Shanghai University
Shanghai 200444, China

Abstract

In this paper, we study the bound on three kinds of hash family using the Singleton bound. To $\varepsilon - U(N; n, m)$ hash family, in the case of $n > m^2 > 1$ and $1 \geq \varepsilon \geq \varepsilon_1(n, m)$, we get that the new bound is better. To $\varepsilon - \Delta U(N; n, m)$ hash family, in the case of $n > m > 1$ and $1 \geq \varepsilon \geq \varepsilon_3(n, m)$, the new bound is better. To $\varepsilon - SU(N; n, m)$ hash family, in the case of $n > 2^m > 2$ and $1 \geq \varepsilon \geq \varepsilon_4(n, m)$, we get that the new bound is better.

Keywords: hash family; the Singleton bound; the Plotkin bound; MDS code

1 Introduction

The concept known as “universal hashing” was invented by Carter and Wegman [1] in 1979. In [2, p. 18], Avi Wigderson characterizes universal hashing as being a tool which “should belong to the fundamental bag of tricks every computer scientist”. The hash function has owned broad use in information authentication field such as digital signature, and has had close relation to authentication codes [10]. In 1980, D. V. Sarwate [5] introduced the Plotkin bound to an $\varepsilon - U(N; n, m)$ hash family, and got $\varepsilon \geq \frac{n-m}{m(n-1)}$. In 1994, D. R. Stinson [3] got $N \geq \frac{n(m-1)}{n(\varepsilon m-1)+m^2(1-\varepsilon)}$ when he studied the $\varepsilon - U(N; n, m)$ hash family. In 1995, he [8] got $N \geq \frac{n(m-1)}{m-n+m\varepsilon(n-1)}$ when studied the $\varepsilon - \Delta U(N; n, m)$ hash family, and got $N \geq 1 + \frac{n(m-1)^2}{m\varepsilon(n-1)+m-n}$ when studied the $\varepsilon - SU(N; n, m)$ hash family.

*Corresponding author: Jiansheng Yang (1963 -), Man, Associate Professor, Doctor, Major in Graph Theory, Coding Theory. E-mail: yjsyjs@staff.shu.edu.cn

In the following, We denote $\frac{(n-m^2)(\log_m n-1)}{(mn-m^2)\log_m n+m^2+n-2mn}$, $\frac{n-m}{m(n-1)}$, $\frac{n(m-1)(m-2)+(\log_2 n+m-1)(n-m)}{m(n-1)(\log_2 n+m-1)-2n(m-1)}$, respectively, by $\varepsilon_1(n, m)$, $\varepsilon_2(n, m)$, $\varepsilon_3(n, m)$ and denote the smaller solution of the equation: $2(n-1)x^2 - [m(n-1)(\log_2 n-3) + 6n-2m-4]x + (m-2)(nm-2n+1) + (n-m)\log_2 n = 0$ by $\varepsilon_4(n, m)$.

In this paper, we introduce the Singleton bound to $\varepsilon - U(N; n, m)$ hash family, and get $N \geq \frac{\log_m n-1}{\varepsilon}$. Through comparing the two bounds, we get that the new bound is better when $1 \geq \varepsilon \geq \varepsilon_1(n, m)$, and the old bound is better when $\varepsilon_1(n, m) > \varepsilon \geq \varepsilon_2(n, m)$. Meanwhile, we introduce the Singleton bound to $\varepsilon - \Delta U(N; n, m)$ hash family, and get $N \geq \frac{\log_2 n+m-1}{m-2+2\varepsilon}$. Through comparing, we get that the new bound is better when $1 \geq \varepsilon \geq \varepsilon_3(n, m)$, and the old bound is better when $\varepsilon_3(n, m) > \varepsilon \geq \frac{1}{m}$. We also introduce the Singleton bound to $\varepsilon - SU(N; n, m)$ hash family, and get $N \geq \frac{m \log_2 n}{m-2(1-\varepsilon)}$. Through comparing, we get that the new bound is better when $1 \geq \varepsilon \geq \varepsilon_4(n, m)$, and the old bound is better when $\varepsilon_4(n, m) > \varepsilon \geq \frac{1}{m}$.

2 Hash Family and Codes

Definition 2.1[7] Let A, B are finite sets, suppose $|A| \geq |B|$, the function $h : A \rightarrow B$ is called hash function.

Definition 2.2[3] Let $\mathcal{h}$ is the set of hash function $h : A \rightarrow B$, if $|A| = n$, $|B| = m$, $|\mathcal{h}| = N$, then it is called $(N; n, m)$ hash family.

Definition 2.3[8] An $(N; n, m)$ hash family is $\varepsilon - universal$ provided that for any two distinct elements $a_1, a_2 \in A$, there exist at most εN functions $h \in \mathcal{h}$ such that $h(a_1) = h(a_2)$. we will use the notation $\varepsilon - U$ as an abbreviation for $\varepsilon - universal$.

If the ε of an $\varepsilon - U(N; n, m)$ hash family is $\frac{1}{m}$, it is known as *universal hashing*[1].

Generally, to an $\varepsilon - U(N; n, m)$ hash family, εN is the smaller the better.

Definition 2.4[8] Suppose that functions in an $(N; n, m)$ hash family, $\mathcal{h}$, have range $B = G$, where G is an additive abelian group (of order m). $\mathcal{h}$ is called $\varepsilon - \Delta universal$ provided that for any two distinct elements $a_1, a_2 \in A$ and for any element $b \in G$, there exist at most εN functions $h \in \mathcal{h}$ such that $h(a_1) - h(a_2) = b$. We will use the notation $\varepsilon - \Delta U$ as an abbreviation for $\varepsilon - \Delta universal$.

Definition 2.5[8] An $(N; n, m)$ hash family is $\varepsilon - strongly universal$ provided that the following two conditions are satisfied:

1. for any element $a \in A$ and any element $b \in B$, there exist exactly N/m functions $h \in \mathcal{h}$ such that $h(a) = b$.
2. for any two distinct elements $a_1, a_2 \in A$ and for any two (not necessarily distinct) elements $b_1, b_2 \in B$, there exist at most $\varepsilon N/m$ functions $h \in \mathcal{h}$ such that $h(a_i) = b_i$, $i = 1, 2$.

We will use the notation ε - SU as an abbreviation for ε -*strongly universal*.

Theorem 2.6[6] If there exists an (N, K, D, q) code, then there exists a $(1 - \frac{D}{N})$ - $U(N; K, q)$ hash family. Conversely, if there exists an ε - $U(N; n, m)$ hash family, then there exists an $(N, n, (1 - \varepsilon)N, m)$ code.

Theorem 2.7[8] If there exists an $[N, k, D, q]$ code C with the property that $e = (1, \dots, 1) \in C$, then there exists a $(1 - \frac{D}{N}) - \Delta U(N; q^{k-1}, q)$ hash family defined over F_q .

Theorem 2.8[5] If there exists an $\varepsilon - U(N; n, m)$ hash family, then

$$\varepsilon \geq \frac{n - m}{m(n - 1)}.$$

Theorem 2.9[8] If there exists an $\varepsilon - \Delta U(N; n, m)$ hash family, then

$$\varepsilon \geq \frac{1}{m}.$$

Theorem 2.10[8] If there exists an $\varepsilon - SU(N; n, m)$ hash family, then

$$\varepsilon \geq \frac{1}{m}.$$

Theorem 2.11[3] If there exists an $\varepsilon - U(N; n, m)$ hash family, then

$$N \geq \frac{n(m - 1)}{n(\varepsilon m - 1) + m^2(1 - \varepsilon)}. \quad (1)$$

Theorem 2.12[8] If there exists an $\varepsilon - \Delta U(N; n, m)$ hash family, then

$$N \geq \frac{n(m - 1)}{m - n + m\varepsilon(n - 1)}. \quad (2)$$

Theorem 2.13[8] If there exists an $\varepsilon - SU(N; n, m)$ hash family, then

$$N \geq 1 + \frac{n(m - 1)^2}{m\varepsilon(n - 1) + m - n}. \quad (3)$$

The following discussion demands $m > 1$.

3 An New Bound for $\varepsilon - U$ Hash Family

Theorem 3.1[4] For $q, n, d \in N$, $q \geq 2$, we have

$$A(n, d) \leq q^{n-d+1}.$$

So, we can get $K \leq q^{N-D+1}$ in an (N, K, D, q) ($q \geq 2$) code. This is called the Singleton bound.

Theorem 3.2 If there exists an $\varepsilon - U(N; n, m)$ hash family, then

$$N \geq \frac{\log_m n - 1}{\varepsilon}. \quad (4)$$

Proof: From Theorem 2.6, since there exists an $\varepsilon - U(N; n, m)$ hash family, then there exists an $(N, n, (1 - \varepsilon)N, m)$ code. Using the Singleton bound, we get

$$n \leq m^{N-(1-\varepsilon)N+1}.$$

So,

$$\log_m n \leq \varepsilon N + 1.$$

Thus,

$$N \geq \frac{\log_m n - 1}{\varepsilon}.$$

□

Lemma 3.3 If $n > m^2$, then

$$1 > \frac{(n - m^2)(\log_m n - 1)}{(mn - m^2) \log_m n + m^2 + n - 2mn} > \frac{n - m}{m(n - 1)}.$$

Proof: Let $\log_m n = 1 + \alpha$, since $n > m^2 > 1$, then $\alpha > 1$. Thus

$$\begin{aligned} & \frac{(n - m^2)(\log_m n - 1)}{(mn - m^2) \log_m n + m^2 + n - 2mn} = \frac{(n - m^2)\alpha}{m(1 + \alpha)(n - m) + m^2 + n - 2mn} \\ & = \frac{(n - m^2)\alpha}{(\alpha - 1)mn + n - m^2\alpha} = \frac{(n - m^2)\alpha}{(\alpha - 1)(m - 1)n + (n - m^2)\alpha} < 1. \end{aligned}$$

Consider function: $f(x) = m^x - mx$ ($m > 1$), $f'(x) = m^x \ln m - m$. Since $m > 1$, then $f'(x) > 0$ when $x > 1$, this means $f(x)$ is a strictly monotony increasing function. Moreover, $f(x) = 0$ when $x = 1$. That is to say, $m^\alpha > m\alpha$ when $n > m^2 > 1$.

So, $n(m - 1)m(-m\alpha + \alpha + m^\alpha - 1) > 0$ is true.

Since $\log_m n = \alpha + 1$ ($\alpha > 1$), then $n = m^{\alpha+1}$, substitute it to the above inequality, we get

$$n(m-1)(-m^2\alpha + m\alpha + n - m) > 0.$$

Now, by this, we have

$$(n\alpha - m^2\alpha)(mn - m) > (mn\alpha - m^2\alpha - mn + n)(n - m).$$

Since

$$mn(\alpha - 1) - m^2\alpha + n = mn(\alpha - 1) - m^2(\alpha - 1) + n - m^2 = m(\alpha - 1)(n - m) + n - m^2 > 0,$$

so,

$$\frac{(n - m^2)\alpha}{mn(\alpha - 1) - m^2\alpha + n} > \frac{n - m}{m(n - 1)}.$$

Thus,

$$\frac{(n - m^2)(\log_m n - 1)}{(mn - m^2)\log_m n + m^2 + n - 2mn} > \frac{n - m}{m(n - 1)}.$$

□

Theorem 3.4 If there exists an $\varepsilon - U(N; n, m)$ hash family, and $n > m^2$, then the bound (4) is better than (1) when $1 \geq \varepsilon \geq \varepsilon_1(n, m)$; the bound (1) is better than (4) when $\varepsilon_1(n, m) > \varepsilon \geq \varepsilon_2(n, m)$.

Proof: The bound (4) better than (1) means

$$\frac{\log_m n - 1}{\varepsilon} \geq \frac{n(m - 1)}{n(\varepsilon m - 1) + m^2(1 - \varepsilon)}.$$

From the above inequality, we have

$$\varepsilon \geq \frac{(n - m^2)(\log_m n - 1)}{(mn - m^2)\log_m n + m^2 + n - 2mn} = \varepsilon_1(n, m).$$

Using the same way, we can get: the bound (1) is better when $\varepsilon_1(n, m) > \varepsilon \geq \varepsilon_2(n, m)$. □

Note: In both bounds, εN must be an integer.

Theorem 3.5 There exists an $\varepsilon - U(N; n, m)$ hash family and $N = \frac{\log_m n - 1}{\varepsilon}$ if and only if there exists an MDS code $(N, n, (1 - \varepsilon)N, m)$.

It is clear. Now, we have:

Theorem 3.6 Suppose q is a power of prime with $(1 < k < n \leq q + 1)$. Then there is a $\frac{k-1}{n} - U(n; q^k, q)$ hash family.

Another nice application also uses MDS code. From [4], we know there exists an $[n, n-1, 2, q]$ ($q \geq 2$) MDS code. Let $n = q^{i+1}$, there exists its subcode $(q^{i+1}, (2q-1) \times q^{q^{i+1}-3}, 2, q)$. Applying Theorem 2.6, the following is obtained.

Theorem 3.7 There exists a $(1 - \frac{2}{q^{i+1}}) - U(q^{i+1}; (2q-1) \times q^{q^{i+1}-3}, q)$ ($q \geq 2, i \geq 1$) hash family.

From this theorem, $m = q, n = (2q-1) \times q^{q^{i+1}-3}, \varepsilon = 1 - \frac{2}{q^{i+1}}$, we have $\varepsilon > \varepsilon_1(n, m)$, then the bound (4) is better. So, $N \geq \lceil \frac{\log_m n - 1}{\varepsilon} \rceil = \lceil [q^{i+1} - 3 - 1 + \log_q(2q-1)] \times \frac{q^{i+1}}{q^{i+1}-2} \rceil = q^{i+1}$. Thus, $(1 - \frac{2}{q^{i+1}}) - U(q^{i+1}; (2q-1) \times q^{q^{i+1}-3}, q)$ hash family has the smallest N .

4 An New Bound for $\varepsilon - \triangle U$ Hash Family

Lemma 4.1 If $n > m$, then

$$1 > \frac{n(m-1)(m-2) + (\log_2 n + m - 1)(n - m)}{m(n-1)(\log_2 n + m - 1) - 2n(m-1)} > \frac{1}{m}.$$

Proof: Since $n > m > 1$, then $(m-1)n(1 - \log_2 n) < 0$. We have

$$n(m-1)(m-2) + (\log_2 n + m - 1)(n - m) < m(n-1)(\log_2 n + m - 1) - 2n(m-1).$$

Since

$$m(n-1)(\log_2 n + m - 1) - 2n(m-1) = m(n-1)(\log_2 n + m - 3) + 2n - 2m > 0,$$

thus,

$$1 > \frac{n(m-1)(m-2) + (\log_2 n + m - 1)(n - m)}{m(n-1)(\log_2 n + m - 1) - 2n(m-1)}.$$

Since

$$m(m-1)(n - \log_2 n) + (m-1)^2(mn - m - 2n) > 0,$$

then

$$mn(m-1)(m-2) + m(\log_2 n + m - 1)(n - m) - [m(n-1)(\log_2 n + m - 1) - 2n(m-1)] > 0.$$

Thus,

$$\frac{n(m-1)(m-2) + (\log_2 n + m - 1)(n - m)}{m(n-1)(\log_2 n + m - 1) - 2n(m-1)} > \frac{1}{m}.$$

□

From the proof of Theorem 2.12 in [8], we have that if there exists an $\varepsilon - \Delta U(N; n, m)$ hash family, then there exists a constant-weight $((N - 1)m, n, 2N(1 - \varepsilon), 2)$ code. Using the Singleton bound, we have

$$n \leq 2^{(N-1)m - 2(1-\varepsilon)N + 1}.$$

So, the bound (4) is changed to

$$N \geq \frac{\log_2 n + m - 1}{m - 2 + 2\varepsilon}. \quad (5)$$

Theorem 4.2 If there exists an $\varepsilon - \Delta U(N; n, m)$ hash family and $n > m$, then the bound (5) is better than (2) when $1 \geq \varepsilon \geq \varepsilon_3(n, m)$; the bound (2) is better than (5) when $\varepsilon_3(n, m) > \varepsilon \geq \frac{1}{m}$.

Proof: The bound (5) better than (2) means

$$\frac{\log_2 n + m - 1}{m - 2 + 2\varepsilon} \geq \frac{n(m-1)}{m - n + m\varepsilon(n-1)}.$$

From the above inequality, we have

$$\varepsilon \geq \frac{n(m-1)(m-2) - (\log_2 n + m - 1)(m - n)}{m(n-1)(\log_2 n + m - 1) - 2n(m-1)} = \varepsilon_3(n, m).$$

Using the same way, we can get: the bound (2) is better when $\varepsilon_3(n, m) > \varepsilon \geq \frac{1}{m}$. □

Note: In both bounds, εN must be an integer.

Example: Let $\varepsilon = 1 - \frac{2}{(q-1)^{i+1}}$ ($q > 2$, $i > 1$), $n = q^{(q-1)^{i+1}-1}$, $m = q$, then through computing, we have $1 > \varepsilon > \varepsilon_3(n, m)$, so the bound (5) is better. Then

$$N \geq \frac{\log_2 n + m - 1}{m - 2 + 2\varepsilon} = \frac{(q-1)^{i+1} \log_2 q + q - \log_2 q - 1}{q + \frac{4}{(q-1)^{i+1}}}.$$

Since $q > 2$, $i > 1$, then we have

$$(q-1)^i < \frac{(q-1)^{i+1} \log_2 q + q - \log_2 q - 1}{q + \frac{4}{(q-1)^{i+1}}} < (q-1)^{i+1}.$$

Since εN is an integer, then $N \geq (q-1)^{i+1}$.

From [4], we know there exists an $[n, n-1, 2, q]$ MDS code C . Let $n = (q-1)^{i+1}$, we may assume that $e = (1, \dots, 1) \in C$, then from Theorem 2.7, we have the following.

Theorem 4.3 There exists a $(1 - \frac{2}{(q-1)^{i+1}}) - \Delta U((q-1)^{i+1}; q^{(q-1)^{i+1}-1}, q)$ ($q > 2$, $i > 1$) hash family.

5 An New Bound for $\varepsilon - SU$ Hash Family

For $n, m > 0$, we denote $2(n-1)$ by a , denote $m(n-1)(\log_2 n - 3) + 6n - 2m - 4$ by b and denote $(m-2)(nm - 2n + 1) + (n-m)\log_2 n$ by c . Then,

$$\varepsilon_4(n, m) = \frac{b - \sqrt{b^2 - 4ac}}{2a}.$$

Lemma 5.1 If $n > 2^m$, then

$$1 > \varepsilon_4(n, m) > \frac{1}{m}.$$

Proof: Since $n > 2^m > 1$, then $n(m-1)(m - \log_2 n) < 0$. So,

$$2(n-1) - [m(n-1)(\log_2 n - 3) + 6n - 2m - 4] + (m-2)(nm - 2n + 1) + (n-m)\log_2 n < 0.$$

This is to say $a + b + c < 0$. Thus,

$$\varepsilon_4(n, m) = \frac{-b - \sqrt{b^2 - 4ac}}{2a} < 1.$$

Since

$$2(n-1) - m[m(n-1)(\log_2 n - 3) + 6n - 2m - 4] + m^2[(m-2)(nm - 2n + 1) + (n-m)\log_2 n] = 2(n-1) + m^2 \log_2 n + m(m^2 - 3m + 4) + 3mn(m-2) + m^3(n - \log_2 n) + m^2 n(m-1)(m-4) > 0.$$

(It is obvious when $m \geq 4$. So, we only need to check on the cases $m = 2$ and $m = 3$ to get the result.)

This is to say,

$$a + bm + cm^2 > 0.$$

Thus,

$$\varepsilon_4(n, m) > \frac{1}{m}.$$

□

From the proof of Theorem 2.13 in [8], we have that if there exists an $\varepsilon - SU(N; n, m)$ hash family, then there exists a constant-weight $(N - 1, n, 2(1 - \varepsilon)N/m, 2)$ code. Using the Singleton bound, we have

$$n \leq 2^{N-1-2N(1-\varepsilon)/m+1}.$$

So, the bound (4) is changed to

$$N \geq \frac{m \log_2 n}{m - 2(1 - \varepsilon)}. \quad (6)$$

Theorem 5.2 If there exists an $\varepsilon - SU(N; n, m)$ hash family and $n > 2^m$, then the bound (6) is better than (3) when $1 \geq \varepsilon \geq \varepsilon_4(n, m)$; the bound (3) is better than (6) when $\varepsilon_4(n, m) > \varepsilon \geq \frac{1}{m}$.

Proof: The bound (6) better than (3) means

$$\frac{m \log_2 n}{m - 2(1 - \varepsilon)} \geq 1 + \frac{n(m - 1)^2}{m\varepsilon(n - 1) + m - n}.$$

From the above inequality, we have

$$1 \geq \varepsilon \geq \varepsilon_4(n, m).$$

Using the same way, we can get: the bound (3) is better when $\varepsilon_4(n, m) > \varepsilon \geq \frac{1}{m}$. □

Note: In both bounds, $\frac{\varepsilon N}{m}$ must be an integer.

References

- [1] J. L. Carter and M. N. Wegman, "Universal Classes of Hash Functions," *Computer and System Sciences*, vol. 18, pp. 143-154, 1979.
- [2] A. Wigderson, "Lectures on the Fusion Method and Eerandomization," Technical Report SOCS-95.2, School of Computer Science, McGill University (file/pub/tech-reports/library/reports/95/TR95.2.ps.gz at the anonymous ftp site ftp.cs.mcgill.ca).

- [3] D. R. Stinson, "Universal Hashing and Authentication Codes," *Designs, Codes and Cryptography*, vol. 4, pp. 369-380, Oct.1994.
- [4] J. H. van Lint, "Introduction to Coding Theory," *Springer-Verlag*, 3rd ed. New York, Inc., pp. 64-69, 1999.
- [5] D. V. Sarwate, "A Note on Universal Classes of Hash Functions," *Information Processing Letters*, vol. 10, pp. 41-45, 1980.
- [6] J. Bierbrauer, T. Johansson, G. Kabatianskii and B. Smeets, "On Families of Hash Functions via Geometric Codes and Concatenation," In "Advances in Cryptology-CRYPTO'93", D. R. Stinson, ed., (Lecture Notes in Computer Science), London: Springer-Verlag, vol. 773, pp. 331-342, 1993.
- [7] Jianzhong ZHANG and GuozhenXIAO, "The Constructions of Authentication Codes With Hashing," *Electronics and Information Technology*, vol.6, pp. 533-539, 2001.
- [8] D. R. Stinson, "On the Connections Between Universal Hashing, Combinatorial Designs and Error-Correcting Codes," *Congressus Numer*, vol. 114, pp. 7-27, 1996.
- [9] D. R. Stinson, "Combinatorial Techniques for Universal Hashing," *Computer and System Sciences*, vol.48, pp. 337-346, Apr. 1994.
- [10] T. Johansson, B. Smeets and G. Kabatianskii, "On the Cardinality of Systematic Authentication Codes via Error Correcting Codes," *IEEE Transactions on Information Theory*, VOL. 42, NO. 2, pp. 566-578, Mar. 1996.